

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.JournalofSurgicalResearch.com

Corrigendum

Corrigendum to: Direct Oral Anticoagulants Are a Potential Alternative to Low-Molecular-Weight Heparin for Thromboprophylaxis in Trauma Patients Sustaining Lower Extremity Fractures [Journal of Surgical Research 258 (2020) 324-331]



Charlie J. Nederpelt, BSc,^{a,b} Kerry A. Breen, BS,^a Majed W. el Hechi, MD,^a Pieta Krijnen, PhD,^b Menno V. Huisman, MD, PhD,^c Inger B. Schipper, MD, PhD,^b Haytham M.A. Kaafarani, MD, MPH,^{a,d} and Martin G. Rosenthal, MD^a

^a Division of Trauma, Emergency Surgery and Surgical Critical Care, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts

^b Department of Trauma Surgery, Leiden University Medical Center, Leiden University, Leiden, the Netherlands

^c Department of Thrombosis and Hemostasis, Leiden University Medical Center, Leiden University, Leiden, the Netherlands

^d Center for Outcomes & Patient Safety in Surgery (COMPASS), Massachusetts General Hospital, Boston, Massachusetts

The authors regret an error in Table 4 stating that the primary outcome of total in-hospital symptomatic venous thromboembolic events occurred in 0.4% of patients receiving low-molecular-weight heparin. The correct symptomatic venous thromboembolism rate in low-molecular-weight heparin

patients in this cohort was 1.4%, equal to the symptomatic venous thromboembolism rate in patients receiving direct oral anticoagulants, as was correctly reported in the text.

The authors would like to apologise for any inconvenience caused.